

## TECHNICAL DATA SHEET

### CyFlow™ Beta2-MG Purified Anti-Hu; Clone B2M-01

**REF** CQ959558

**For Research Use Only.  
Not for use in diagnostic or therapeutic procedures.**

### Specifications

<b>Antigen</b>	β2-microglobulin
<b>Alternative Names</b>	—
<b>Clone</b>	B2M-01
<b>Clonality</b>	monoclonal
<b>Format</b>	Purified
<b>Host / Isotype</b>	Mouse / IgG2a
<b>Species Reactivity</b>	Human
<b>Negative Species Reactivity</b>	Mouse   Cow   Dog   Rabbit   Chicken
<b>Quantity [Concentration]</b>	0.1 mg [ 1 mg/ml ]
<b>Immunogen</b>	Purified human β2-microglobulin

### Specificity

The mouse monoclonal antibody B2M-01 recognizes β2-microglobulin associated with cell-surface MHC Class I molecules and other membrane antigens as well as with soluble β2-microglobulin. β2-

#### Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany  
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: [info@sysmex-partec.com](mailto:info@sysmex-partec.com)

microglobulin is a 12 kDa Ig like glycoprotein expressed on lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells and epithelial cells. It is absent on erythrocytes.

## Application

Based on published sources, this antibody is suitable for the following applications:

- Flow cytometry
- Immunoprecipitation
- Western blot
- Immunohistochemistry (paraffin-embedded sections)
- Immunocytochemistry
- Enzyme-linked immunosorbent assay
- Radioimmunoassay

## Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH  $\approx$ 7.4, containing 0.1% (w/v) sodium azide.

## Storage and Stability

<b>Storage</b>	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
<b>Stability</b>	Do not use after expiration date stamped on vial label.

## Background Information

$\beta$ 2-microglobulin ( $\beta$ 2M) non-covalently associates with the 44 kDa  $\alpha$  chain to form the HLA Class I antigen complex. Human  $\beta$ 2-microglobulin associated with HLA Class I antigens is expressed on many types of cells including lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells, and epithelial cells. It is absent on erythrocytes.

## References

- Hilgert I, Horejsi V, Kristofova H: The use of murine monoclonal antibody B2M-01 for detection and purification of human beta 2-microglobulin. Folia Biol (Praha). 1984; 30(6):369-76. < PMID: 6394369 >
- Lafon M, Prehaud C, Megret F, Lafage M, Mouillot G, Roa M, Moreau P, Rouas-Freiss N, Carosella ED: Modulation of HLA-G Expression in Human Neural Cells after Neurotropic Viral Infections. J Virol. 2005 Dec; 79(24):15226-15237. < PMID: 16306594 >

---

### Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany  
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: [info@sysmex-partec.com](mailto:info@sysmex-partec.com)

- Khurana M, Traum AZ, Aivado M, Wells MP, Guerrero M, Grall F, Libermann TA, Schachter AD: Urine proteomic profiling of pediatric nephrotic syndrome. *Pediatr Nephrol.* 2006 Sep; 21(9):1257-65. < PMID: 16810512 >
- Poláková K, Železníková T, Russ G: HLA-G5 in the blood of leukemia patients and healthy individuals. *Leuk Res.* 2013 Feb; 37(2):139-45. < PMID: 23146557 >

---

The Safety Data Sheet for this product is available at [www.sysmex-partec.com/services](http://www.sysmex-partec.com/services).

---

---

**Contact Information:**

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany  
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: [info@sysmex-partec.com](mailto:info@sysmex-partec.com)