

TECHNICAL DATA SHEET

CyFlow™ CD4 Azide Free Anti-Ms; Clone GK1.5

REF CN672458

**For Research Use Only.
Not for use in diagnostic or therapeutic procedures.**

Specifications

Antigen	CD4
Alternative Names	T4, Leu3a
Clone	GK1.5
Clonality	monoclonal
Format	Azide Free
Host / Isotype	Rat / IgG2b
Species Reactivity	Mouse
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	Mouse CTL clone V4 cells

Specificity

The rat monoclonal antibody GK1.5 recognizes an extracellular epitope of mouse CD4 transmembrane glycoprotein (55 kDa).

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

Based on published sources, this antibody is suitable for the following applications:

- Flow cytometry
- Immunoprecipitation
- Immunohistochemistry (frozen sections)
- Immunocytochemistry
- Functional assays

Storage Buffer

The reagent is provided in azide-free phosphate buffered saline (PBS) solution, pH \approx 7.4; 0.2 μ m filter sterilized.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD4 is a single chain transmembrane glycoprotein of immunoglobulin supergene family. In its extracellular region there are 4 immunoglobulin-like domains (1 Ig-like V-type and 3 Ig-like C2-type). The intracellular region of CD4 associates with p56Lck, a Src-like protein tyrosine kinase. It was described that CD4 segregates into specific detergent-resistant T-cell membrane microdomains. CD4 binds to MHC class II molecules (by CDR2-like region in CD4 domain 1), HIV envelope protein gp120 (by CDR2-like region in CD4 domain 1) and other ligands, such as IL-16 (by to CD4 domain 3) or L-selectin. CD4 is a co-receptor involved in immune response (co-receptor activity in binding to MHC class II molecules) and HIV infection. CD4 regulates T-cell activation, T/B-cell adhesion, T-cell differentiation, T-cell selection and signal transduction. Defects in antigen presentation (MHC class II) cause dysfunction of CD4+ T-cells and their almost complete absence in patients blood, tissue and organs (SCID immunodeficiency).

References

- Dialynas DP, Wilde DB, Marrack P, Pierres A, Wall KA, Havran W, Otten G, Loken MR, Pierres M, Kappler J, et al.: Characterization of the murine antigenic determinant, designated L3T4a, recognized by monoclonal antibody GK1.5: expression of L3T4a by functional T cell clones appears to correlate primarily with class II MHC antigen-reactivity. Immunol Rev. 1983; 74:29-56. < PMID: 6195085 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Dialynas DP, Quan ZS, Wall KA, Pierres A, Quintáns J, Loken MR, Pierres M, Fitch FW: Characterization of the murine T cell surface molecule, designated L3T4, identified by monoclonal antibody GK1.5: similarity of L3T4 to the human Leu-3/T4 molecule. *J Immunol.* 1983 Nov; 131(5):2445-51. < PMID: 6415170 >
- Wilde DB, Marrack P, Kappler J, Dialynas DP, Fitch FW: Evidence implicating L3T4 in class II MHC antigen reactivity: monoclonal antibody GK1.5 (anti-L3T4a) blocks class II MHC antigen-specific proliferation, release of lymphokines, and binding by cloned murine helper T lymphocyte lines. *J Immunol.* 1983 Nov; 131(5):2178-83. < PMID: 6195255 >
- Wu L, Antica M, Johnson GR, Scollay R, Shortman K: Developmental potential of the earliest precursor cells from the adult mouse thymus. *J Exp Med.* 1991 Dec 1; 174(6):1617-27. < PMID: 1683894 >
- Godfrey DI, Kennedy J, Gately MK, Hakimi J, Hubbard BR, Zlotnik A: IL-12 influences intrathymic T cell development. *J Immunol.* 1994 Mar 15; 152(6):2729-35. < PMID: 7511624 >
- Gavett SH, Chen X, Finkelman F, Wills-Karp M: Depletion of murine CD4+ T lymphocytes prevents antigen-induced airway hyperreactivity and pulmonary eosinophilia. *Am J Respir Cell Mol Biol.* 1994 Jun; 10(6):587-93. < PMID: 8003337 >
- Zheng B, Han S, Kelsoe G: T helper cells in murine germinal centers are antigen-specific emigrants that downregulate Thy-1. *J Exp Med.* 1996 Sep 1; 184(3):1083-91. < PMID: 9064325 >
- Felix NJ, Donermeyer DL, Horvath S, Walters JJ, Gross ML, Suri A, Allen PM: Alloreactive T cells respond specifically to multiple distinct peptide-MHC complexes. *Nat Immunol.* 2007 Apr; 8(4):388-97. < PMID: 17322886 >
- Yi H, Zhen Y, Zeng C, Zhang L, Zhao Y: Depleting anti-CD4 monoclonal antibody (GK1.5) treatment: influence on regulatory CD4+CD25+Foxp3+ T cells in mice. *Transplantation.* 2008 Apr 27; 85(8):1167-74. < PMID: 18431238 >
- Hu M, Watson D, Zhang GY, Graf N, Wang YM, Sartor M, Howden B, Fletcher J, Alexander SI: Long-term cardiac allograft survival across an MHC mismatch after "pruning" of alloreactive CD4 T cells. *J Immunol.* 2008 May 15; 180(10):6593-603. < PMID: 18453578 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com