

TECHNICAL DATA SHEET

CyFlow™ IFN-gamma Purified Anti-Hu; Clone 4S.B3

REF AJ218636

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	IFN- γ
Alternative Names	—
Clone	4S.B3
Clonality	monoclonal
Format	Purified
Host / Isotype	Mouse / IgG1
Species Reactivity	Human, Non-Human Primates
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	Interferon gamma derived from human leukocytes

Specificity

The mouse monoclonal antibody 4S.B3 recognizes IFN- γ antigen, a 16-25 kDa cytokine produced by activated Th1 cells and NK cells. Binds both glycosylated and non-glycosylated protein.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

Based on published sources, this antibody is suitable for the following applications:

- Flow cytometry
- Immunoprecipitation
- Western blot
- Immunohistochemistry (paraffin-embedded sections)
- Immunocytochemistry
- Functional assays
- Enzyme-linked immunosorbent assay
- Radioimmunoassay

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH \approx 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

The Interferon γ (IFN- γ) is an important regulator of the immune response, produced in activated Th1 cells and NK cells, particularly in response to IL-2, TNF- α and IL-12; its production is suppressed by IL-4, IL-10, and TGF- β . The production of IFN- γ is activated by specific antigens or mitogens through the T cell antigen receptor. IFN- γ polypeptide forms: 40-60 kDa forms are observable under non-denaturing conditions as dimers and trimers; 20 kDa and 25 kDa forms exist due to variable glycosylation. IFN- γ belongs to the type II interferons, also called immune IFN. IFN- γ shows antiviral activity and has important immunoregulatory functions. It is a potent activator of macrophages and had antiproliferative effects on transformed cells. IFN- γ plays an important role in regulating B cell differentiation by simultaneously stimulating class switch recombination to the IgG3 and IgG2a isotypes while repressing class switch recombination to the IgE and IgG1 isotypes. It also appears to promote antigen presentation by B cells through its effects on MHC. Binding of IFN- γ to its receptor increases the expression of class I MHC on all somatic cells. It also enhances the expression of class II MHC on antigen-presenting cells. IFN- γ is the major means by which T cells activate macrophages, increasing their ability to kill bacteria, parasites,

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

and tumors. The activation of macrophages by IFN- γ is essential for the elimination of bacteria that replicate within the phagosomes of macrophages (f.e. Mycobacteria and *Listeria monocytogenes*). IFN- γ can potentiate the high antiviral and antitumor effects of the type I interferons (IFN- α , IFN- β). IFN- γ may also activate neutrophils and NK cells.

References

- Meager A, Parti S, Barwick S, Spragg J, O'Hagan K: Detection of hybridomas secreting monoclonal antibodies to human gamma interferon using a rapid screening technique and specificity of certain monoclonal antibodies to gamma interferon. *J Interferon Res.* 1984 Fall; 4(4):619-25. < PMID: 6438252 >
- Caulfield JJ, Fernandez MH, Sousa AR, Lane SJ, Lee TH, Hawrylowicz CM: Regulation of major histocompatibility complex class II antigens on human alveolar macrophages by granulocyte-macrophage colony-stimulating factor in the presence of glucocorticoids. *Immunology.* 1999 Sep; 98(1):104-10. < PMID: 10469240 >
- Coles AJ, Wing M, Smith S, Coraddu F, Greer S, Taylor C, Weetman A, Hale G, Chatterjee VK, Waldmann H, Compston A: Pulsed monoclonal antibody treatment and autoimmune thyroid disease in multiple sclerosis. *Lancet.* 1999 Nov 13; 354(9191):1691-5. < PMID: 10568572 >
- Brattig NW, Lepping B, Timmann C, Büttner DW, Marfo Y, Hamelmann C, Horstmann RD: *Onchocerca volvulus*-exposed persons fail to produce interferon-gamma in response to *O. volvulus* antigen but mount proliferative responses with interleukin-5 and IL-13 production that decrease with increasing microfilarial density. *J Infect Dis.* 2002 Apr 15; 185(8):1148-54. < PMID: 11930325 >
- Janke M, Witsch EJ, Mages HW, Hutloff A, Kroczeck RA: Eminent role of ICOS costimulation for T cells interacting with plasmacytoid dendritic cells. *Immunology.* 2006 Jul; 118(3):353-60. < PMID: 16827896 >
- Kosub DA, Lehrman G, Milush JM, Zhou D, Chacko E, Leone A, Gordon S, Silvestri G, Else JG, Keiser P, Jain MK, Sodora DL: Gamma/Delta T-Cell Functional Responses Differ after Pathogenic Human Immunodeficiency Virus and Nonpathogenic Simian Immunodeficiency Virus Infections. *J Virol.* 2008 Feb; 82(3):1155-1165. < PMID: 18045946 >
- Erik L Brincks, Tamara A Kucaba, Kevin L Legge, Thomas S Griffith: Influenza-induced expression of functional TNF-related apoptosis-inducing ligand (TRAIL) on human PBMC. *Hum Immunol.* 2008 Oct; 69(10):634-646. < PMID: 18723061 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com