

TECHNICAL DATA SHEET

CyFlow™ CD140a Biotin Anti-Hu; Clone 16A1

REF AS196282

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD140a
Alternative Names	PDGFRA, PDGFRF2
Clone	16A1
Clonality	monoclonal
Format	Biotin
Host / Isotype	Mouse / IgG1
Species Reactivity	Human
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	CD140a-transfected NIH 3T3 cells

Specificity

The mouse monoclonal antibody 16A1 recognizes CD140a antigen, the 170 kDa α chain of platelet-derived growth factor receptor, which is widely expressed on a variety of mesenchymal-derived cells and plays pro-proliferative or anti-proliferative roles in various tumors.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for indirect immunofluorescence analysis by Flow Cytometry. Suggested working usage is 3 µg/ml. Indicated dilution is recommended starting point for use of this product, but working concentrations should be validated by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD140a (PDGF-RA; platelet-derived growth factor receptor α) is a cell surface receptor for members of platelet-derived growth factor family, whose intracellular part contains a tyrosine kinase domain. CD140a forms homodimers, or heterodimerizes with CD140b (PDGF-RB). Whereas CD140b induces in different cell types their proliferation and migration, the role of CD140a is more controversial, with pro-proliferative or anti-proliferative effects. CD140a has early developmental functions, mediates mesodermal cell migration, and later acts in signaling associated in epithelial-mesenchymal interactions.

References

- Kish DD, Gorbachev AV, Fairchild RL: CD8+ T cells produce IL-2, which is required for CD(4+)CD25+ T cell regulation of effector CD8+ T cell development for contact hypersensitivity responses. J Leukoc Biol. 2005 Sep; 78(3):725-35. < PMID: 16000396 >
- Schmahl J, Rizzolo K, Soriano P: The PDGF signaling pathway controls multiple steroid-producing lineages. Genes Dev. 2008 Dec 1; 22(23):3255-67. < PMID: 19056881 >
- Andrae J, Gallini R, Betsholtz C: Role of platelet-derived growth factors in physiology and medicine. Genes Dev. 2008 May 15; 22(10):1276-312. < PMID: 18483217 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- French WJ, Creemers EE, Tallquist MD: Platelet-derived growth factor receptors direct vascular development independent of vascular smooth muscle cell function. Mol Cell Biol. 2008 Sep; 28(18):5646-57. < PMID: 18606782 >
- Faraone D, Aguzzi MS, Toietta G, Facchiano AM, Facchiano F, Magenta A, Martelli F, Truffa S, Cesareo E, Ribatti D, Capogrossi MC, Facchiano A: Platelet-derived growth factor-receptor alpha strongly inhibits melanoma growth in vitro and in vivo. Neoplasia. 2009 Aug; 11(8):732-42. < PMID: 19649203 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com