

## TECHNICAL DATA SHEET

### **CyFlow™ CD158d PE** **Anti-Hu; Clone mAb#33**

**REF** AG800254

**For Research Use Only.**

**Not for use in diagnostic or therapeutic procedures.**

### Specifications

<b>Antigen</b>	CD158d
<b>Alternative Names</b>	KIR2DL4, KIR103, 103AS, 15.212, KIR103AS
<b>Clone</b>	mAb#33
<b>Clonality</b>	monoclonal
<b>Format</b>	PE
<b>Host / Isotype</b>	Mouse / IgG1
<b>Species Reactivity</b>	Human
<b>Negative Species Reactivity</b>	—
<b>Quantity</b>	100 tests
<b>Immunogen</b>	NK3.3 cells and KIR2DL4-Ig fusion protein

### Specificity

The mouse monoclonal antibody mAb#33 (also known as mAb 33 or 33) recognizes extracellular portion of CD158d antigen, a 45 kDa NK cell marker. Cell surface expression and function of CD158d depends on genotype of particular individuals.

#### Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany  
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: [info@sysmex-partec.com](mailto:info@sysmex-partec.com)

## Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 20 µl reagent / 100 µl of whole blood or 10<sup>6</sup> cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

## Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

## Storage and Stability

<b>Storage</b>	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
<b>Stability</b>	Do not use after expiration date stamped on vial label.

## Background Information

CD158d (KIR2DL4) is a KIR family member that shares structural features with both activating and inhibitory receptors and may mediate different functions under different circumstances. It contains cytoplasmic ITIM, suggesting inhibitory function, but also transmembrane domain similar to those of activating KIRs. It has been reported that CD158d serves as an inhibitory receptor for peripheral and uterine NK cells, but its ligation with soluble mAbs (unlike immobilized mAbs) results in activation of IFN-γ secretion. CD158d also binds both membrane form and soluble form of its ligand HLA-G.

## References

- Rajagopalan S, Fu J, Long EO: Cutting edge: induction of IFN-gamma production but not cytotoxicity by the killer cell Ig-like receptor KIR2DL4 (CD158d) in resting NK cells. J Immunol. 2001 Aug 15; 167(4):1877-81. < PMID: 11489965 >
- Goodridge JP, Witt CS, Christiansen FT, Warren HS: KIR2DL4 (CD158d) genotype influences expression and function in NK cells. J Immunol. 2003 Aug 15; 171(4):1768-74. < PMID: 12902476 >
- Yan WH, Fan LA: Residues Met76 and Gln79 in HLA-G alpha1 domain involve in KIR2DL4 recognition. Cell Res. 2005 Mar; 15(3):176-82. < PMID: 15780179 >

---

### Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany  
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: [info@sysmex-partec.com](mailto:info@sysmex-partec.com)

- LeMaout J, Zafaranloo K, Le Danff C, Carosella ED: HLA-G up-regulates ILT2, ILT3, ILT4, and KIR2DL4 in antigen presenting cells, NK cells, and T cells. FASEB J. 2005 Apr; 19(6):662-4. < PMID: 15670976 >
- Rajagopalan S, Bryceson YT, Kuppusamy SP, Geraghty DE, van der Meer A, Joosten I, Long EO: Activation of NK cells by an endocytosed receptor for soluble HLA-G. PLoS Biol. 2006 Jan; 4(1):e9. < PMID: 16366734 >

---

The Safety Data Sheet for this product is available at [www.sysmex-partec.com/services](http://www.sysmex-partec.com/services).

---

---

**Contact Information:**

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany  
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: [info@sysmex-partec.com](mailto:info@sysmex-partec.com)