

## TECHNICAL DATA SHEET

### CyFlow™ CD163 APC Anti-Hu; Clone GHI/61

**REF** AH476333

---

**For Research Use Only.  
Not for use in diagnostic or therapeutic procedures.**

---

## Specifications

<b>Antigen</b>	CD163
<b>Alternative Names</b>	GH1/61, D11, RM3/1, M130
<b>Clone</b>	GHI/61
<b>Clonality</b>	monoclonal
<b>Format</b>	APC
<b>Host / Isotype</b>	Mouse / IgG1
<b>Species Reactivity</b>	Human
<b>Negative Species Reactivity</b>	—
<b>Quantity</b>	100 tests
<b>Immunogen</b>	Hairy cell leukemia cells

## Specificity

The mouse monoclonal antibody GHI/61 recognizes CD163 antigen, a 130 kDa high affinity scavenger receptor expressed mainly on monocytes and macrophages, which binds hemoglobin-haptoglobin complex.

---

**Contact Information:**

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany  
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: [info@sysmex-partec.com](mailto:info@sysmex-partec.com)

## Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 10 µl reagent / 100 µl of whole blood or 10<sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

## Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

## Storage and Stability

<b>Storage</b>	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
<b>Stability</b>	Do not use after expiration date stamped on vial label.

## Background Information

CD163 (M130) is a member of the scavenger receptor family, accounting for the clearance of hemoglobin-haptoglobin complexes during limited hemolysis, which protects the body, in particular the kidneys, against heme-mediated oxidative damages. It does not have measurable affinity for noncomplexed hemoglobin or haptoglobin. Immunomodulatory role of CD163 has been postulated. CD163 is expressed by cells of the monocyte-macrophage lineage and its extracellular part also circulates in plasma as a soluble protein, especially during sepsis and other conditions affecting macrophage activity, when its level may raise manifold.

## References

- Pulford K, Micklem K, McCarthy S, Cordell J, Jones M, Mason DY: A monocyte/macrophage antigen recognized by the four antibodies GHI/61, Ber-MAC3, Ki-M8 and SM4. *Immunology*. 1992 Apr; 75(4):588-95. < PMID: 1592433 >
- Law SK, Micklem KJ, Shaw JM, Zhang XP, Dong Y, Willis AC, Mason DY: A new macrophage differentiation antigen which is a member of the scavenger receptor superfamily. *Eur J Immunol*. 1993 Sep; 23(9):2320-5. < PMID: 8370408 >

---

### Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany  
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: [info@sysmex-partec.com](mailto:info@sysmex-partec.com)

- Kishimoto T, Goyert S, Kikutani H, Mason D, Miyasaka M, Moretta L, Ohno T, Okumura K, Shaw S, Springer TA, Sugamura K, Sugawara H, von dem Borne AEGK, Zola H (Eds): Leucocyte Typing VI. Garland Publishing Inc, New York. 1997; 1-1342. < NLM ID: 9712219 >
- Møller HJ, Peterslund NA, Graversen JH, Moestrup SK: Identification of the hemoglobin scavenger receptor/CD163 as a natural soluble protein in plasma. *Blood*. 2002 Jan 1; 99(1):378-80. < PMID: 11756196 >
- Madsen M, Møller HJ, Nielsen MJ, Jacobsen C, Graversen JH, van den Berg T, Moestrup SK: Molecular characterization of the haptoglobin-hemoglobin receptor CD163: Ligand binding properties of the scavenger receptor cysteine-rich domain region. *J Biol Chem*. 2004 Dec 3; 279(49):51561-7. < PMID: 15448162 >
- Philippidis P, Mason JC, Evans BJ, Nadra I, Taylor KM, Haskard DO, Landis RC: Hemoglobin scavenger receptor CD163 mediates interleukin-10 release and heme oxygenase-1 synthesis: antiinflammatory monocyte-macrophage responses in vitro, in resolving skin blisters in vivo, and after cardiopulmonary bypass surgery. *Circ Res*. 2004 Jan 9; 94(1):119-26. < PMID: 14656926 >
- Kim WK, Alvarez X, Fisher J, Bronfin B, Westmoreland S, McLaurin J, Williams K: CD163 identifies perivascular macrophages in normal and viral encephalitic brains and potential precursors to perivascular macrophages in blood. *Am J Pathol*. 2006 Mar; 168(3):822-34. < PMID: 16507898 >
- Moniuszko M, Kowal K, Rusak M, Pietruczuk M, Dabrowska M, Bodzenta-Lukaszyk A: Monocyte CD163 and CD36 expression in human whole blood and isolated mononuclear cell samples: influence of different anticoagulants. *Clin Vaccine Immunol*. 2006 Jun; 13(6):704-7. < PMID: 16760331 >
- Bover LC, Cardó-Vila M, Kuniyasu A, Sun J, Rangel R, Takeya M, Aggarwal BB, Arap W, Pasqualini R: A previously unrecognized protein-protein interaction between TWEAK and CD163: potential biological implications. *J Immunol*. 2007 Jun 15; 178(12):8183-94. < PMID: 17548657 >
- Kusi KA, Gyan BA, Goka BQ, Dodoo D, Obeng-Adjei G, Troye-Blomberg M, Akanmori BD, Adjimani JP: Levels of soluble CD163 and severity of malaria in children in Ghana. *Clin Vaccine Immunol*. 2008 Sep; 15(9):1456-60. < PMID: 18632918 >

---

The Safety Data Sheet for this product is available at [www.sysmex-partec.com/services](http://www.sysmex-partec.com/services).

---

---

**Contact Information:**

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany  
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: [info@sysmex-partec.com](mailto:info@sysmex-partec.com)