

TECHNICAL DATA SHEET

CyFlow™ CD62P Biotin Anti-Hu; Clone AK4

REF CU508361

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

| | |
|------------------------------------|---------------------------|
| Antigen | CD62P |
| Alternative Names | P-selectin |
| Clone | AK4 |
| Clonality | monoclonal |
| Format | Biotin |
| Host / Isotype | Mouse / IgG1 |
| Species Reactivity | Human, Non-Human Primates |
| Negative Species Reactivity | — |
| Quantity [Concentration] | 0.1 mg [1 mg/ml] |
| Immunogen | Human platelets |

Specificity

The mouse monoclonal antibody AK4 recognizes CD62P antigen, a 140 kD single chain type I transmembrane glycoprotein present in secretory α -granules in platelets, in Weibel-Palade bodies in endothelial cells and in megakaryocytes; it is relocated to the plasma membrane upon activation.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for indirect immunofluorescence analysis by Flow Cytometry. Working concentrations should be determined by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH \approx 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

| | |
|------------------|---|
| Storage | Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze. |
| Stability | Do not use after expiration date stamped on vial label. |

Background Information

CD62P (P-selectin) is an adhesion glycoprotein that is expressed on platelets and endothelial cells upon their activation. Interaction between CD62P and its mucin-like ligand PSGL-1 (P-selectin glycoprotein ligand-1) expressed on the microvilli of most leukocytes supports leukocyte rolling along postcapillary venules at the earliest time of inflammation. Both CD62P and PSGL-1 are extended glycoproteins that form homodimers. CD62P dimerization is probably mediated through interactions of the transmembrane domains and stabilizes leukocyte tethering and rolling, probably by increasing rebinding within a bond cluster.

References

- Dunlop LC, Skinner MP, Bendall LJ, Favalaro EJ, Castaldi PA, Gorman JJ, Gamble JR, Vadas MA, Berndt MC: Characterization of GMP-140 (P-selectin) as a circulating plasma protein. J Exp Med. 1992 Apr 1; 175(4):1147-50. < PMID: 1372646 >
- Holme PA, Müller F, Solum NO, Brosstad F, Frøland SS, Aukrust P: Enhanced activation of platelets with abnormal release of RANTES in human immunodeficiency virus type 1 infection. FASEB J. 1998 Jan; 12(1):79-89. < PMID: 9438413 >
- Kowalska MA, Ratajczak J, Hoxie J, Brass LF, Gewirtz A, Poncz M, Ratajczak MZ: Megakaryocyte precursors, megakaryocytes and platelets express the HIV co-receptor CXCR4 on their surface:

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

determination of response to stromal-derived factor-1 by megakaryocytes and platelets. Br J Haematol. 1999 Feb; 104(2):220-9. < PMID: 10050701 >

- Ludwig RJ, Schultz JE, Boehncke WH, Podda M, Tandi C, Krombach F, Baatz H, Kaufmann R, von Andrian UH, Zollner TM: Activated, not resting, platelets increase leukocyte rolling in murine skin utilizing a distinct set of adhesion molecules. J Invest Dermatol. 2004 Mar; 122(3):830-6. < PMID: 15086572 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com