

TECHNICAL DATA SHEET

CyFlow™ CD8 Purified Anti-Hu; Clone MEM-87

REF CX396178

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD8
Alternative Names	—
Clone	MEM-87
Clonality	monoclonal
Format	Purified
Host / Isotype	Mouse / IgG1
Species Reactivity	Human
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	Leukocytes of a patient suffering from LGL-type Leukemia

Specificity

The mouse monoclonal antibody MEM-87 recognizes CD8 antigen, a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. CD8

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

is a disulfide-linked dimer and exists as a CD8 α/α homodimer or CD8 α/β heterodimer (each monomer approx. 32-34 kDa).

Application

Based on published sources, this antibody is suitable for the following applications:

- Flow cytometry
- Immunoprecipitation

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH \approx 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

The CD8 T cell coreceptor (monomer approx. 32-34 kDa) is expressed as α/β heterodimer on majority of MHC I-restricted conventional T cells and thymocytes and as α/α homodimer on subsets of memory T cells, intraepithelial lymphocytes, NK cells and dendritic cells. Regulation of CD8 β level on T cell surface seems to be an important mechanism to control their effector function. Assembly of CD8 α/β but not α/α dimers is connected with formation or localization to the lipid rafts. Recruiting triggered TCR complexes to these membrane microdomains as well as affinity of TCR to MHC I is modulated by CD8, thereby affecting the functional diversity of the TCR signaling.

References

- Schlossman SF, Boumsell L, Gilks W, Harlan JM, Kishimoto T, Morimoto C, Ritz J, Shaw S, Silverstein R, Springer T, Tedder T, Todd R (Eds): Leucocyte Typing V. Oxford University Press, Oxford. 1995; < NLM ID: 9514353 >
- Devine L, Thakral D, Nag S, Dobbins J, Hodsdon ME, Kavathas PB: Mapping the binding site on CD8 beta for MHC class I reveals mutants with enhanced binding. J Immunol. 2006 Sep 15; 177(6):3930-8. < PMID: 16951356 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Pang DJ, Hayday AC, Bijlmakers MJ: CD8 Raft localization is induced by its assembly into CD8alpha beta heterodimers, Not CD8alpha alpha homodimers. J Biol Chem. 2007 May 4; 282(18):13884-94. < PMID: 17341584 >
- van den Berg HA, Wooldridge L, Laugel B, Sewell AK: Coreceptor CD8-driven modulation of T cell antigen receptor specificity. J Theor Biol. 2007 Nov 21; 249(2):395-408. < PMID: 17869274 >
- Supper V, Schiller HB, Paster W, Forster F, Boulègue C, Mitulovic G, Leksa V, Ohradanova-Repic A, Machacek C, Schatzlmaier P, Zlabinger GJ, Stockinger H: Association of CD147 and Calcium Exporter PMCA4 Uncouples IL-2 Expression from Early TCR Signaling. J Immunol. 2016 Feb 1; 196(3):1387-99. < PMID: 26729804 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com