

TECHNICAL DATA SHEET

CyFlow™ CD82 Purified Anti-Hu; Clone C33

REF CS166770

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD82
Alternative Names	R2I 4F9, C33
Clone	C33
Clonality	monoclonal
Format	Purified
Host / Isotype	Mouse / IgG2a
Species Reactivity	Human
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	C91/PL (human HTLV-1+ T cell line)

Specificity

The mouse monoclonal antibody C33 recognizes CD82 antigen, a widely expressed cell surface protein of the tetraspanin family. CD82 is also found in endosome/lysosome compartments.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

Based on published sources, this antibody is suitable for the following applications:

- Flow cytometry
- Immunoprecipitation
- Western blot
- Immunohistochemistry (paraffin-embedded sections)
- Immunocytochemistry
- Functional assays

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH \approx 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD82 (KAI1), a member of the tetraspanin family, forms complexes with other tetraspanin proteins, integrins, coreceptors, MHC class I and II molecules. These complexes influence adhesion, morphology, activation, proliferation and differentiation of B, T and other cells. CD82 regulates cytoskeleton rearrangement and may participate in the turnover of the tetraspanin complex members. Besides in the plasma membrane, CD82 is localized also in endosome/lysosome compartments. Tumor-suppressive roles of CD82 have been demonstrated.

References

- Fukudome K, Furuse M, Imai T, Nishimura M, Takagi S, Hinuma Y, Yoshie O: Identification of membrane antigen C33 recognized by monoclonal antibodies inhibitory to human T-cell leukemia virus type 1 (HTLV-1)-induced syncytium formation: altered glycosylation of C33 antigen in HTLV-1-positive T cells. J Virol. 1992 Mar; 66(3):1394-401. < PMID: 1738199 >
- Imai T, Yoshie O: C33 antigen and M38 antigen recognized by monoclonal antibodies inhibitory to syncytium formation by human T cell leukemia virus type 1 are both members of the transmembrane 4

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- superfamily and associate with each other and with CD4 or CD8 in T cells. J Immunol. 1993 Dec 1; 151(11):6470-81. < PMID: 8245480 >
- Imai T, Kakizaki M, Nishimura M, Yoshie O: Molecular analyses of the association of CD4 with two members of the transmembrane 4 superfamily, CD81 and CD82. J Immunol. 1995 Aug 1; 155(3):1229-39. < PMID: 7636191 >
 - Ueda T, Ichikawa T, Tamaru J, Mikata A, Akakura K, Akimoto S, Imai T, Yoshie O, Shiraishi T, Yatani R, Ito H, Shimazaki J: Expression of the KAI1 protein in benign prostatic hyperplasia and prostate cancer. Am J Pathol. 1996 Nov; 149(5):1435-40. < PMID: 8909232 >
 - Escola JM, Kleijmeer MJ, Stoorvogel W, Griffith JM, Yoshie O, Geuze HJ: Selective enrichment of tetraspan proteins on the internal vesicles of multivesicular endosomes and on exosomes secreted by human B-lymphocytes. J Biol Chem. 1998 Aug 7; 273(32):20121-7. < PMID: 9685355 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com